

The Sum Rules for Structure Functions of Polarized $e(\mu)N$ Scattering: Theory vs. Experiment

B.L.Ioffe

Institute of Theoretical and Experimental Physics
117259, Moscow, Russia

Abstract

The determination of twist-4 corrections to the structure functions of polarized $e(\mu)N$ scattering by QCD sum rules is reviewed and critically analyzed. It is found that in the case of the Bjorken sum rule the twist-4 correction is small at $Q^2 > 5 \text{ GeV}^2$. However, the accuracy of the today experimental data is insufficient to reliably determine α_s from the Bjorken sum rule. For the singlet sum rule – $p+n$ – the QCD sum rule gives only the order of magnitude of twist-4 correction. At low and intermediate Q^2 the model is presented which realizes a smooth connection of the Gerasimov-Drell-Hearn sum rules at $Q^2 = 0$ with the sum rules for $\Gamma_{p,n}(Q^2)$ at high Q^2 . The model is in a good agreement with the experiment.

1. Introduction

In the last few years there is a strong interest to the problem of nucleon spin structure: how nucleon spin is distributed among its constituents - quarks and gluons. New experimental data continuously appear and precision increases (for the recent data see [1], [2]). One of the most important item of the information comes from the measurements of the first moment of the spin-dependent nucleon structure functions $g_1(x)$ which determine the parts of nucleon spin carried by u , d and s quarks and gluons. The accuracy of the data is now of a sort that the account of twist-4 terms is of importance when comparing the data with the Bjorken and Ellis-Jaffe sum rules at high Q^2 . On the other side, at low and intermediate Q^2 a smooth connection of the sum rules for the first moments of $g_1(x, Q^2)$ with the Gerasimov-Drell-Hearn (GDH) sum rules [3,4] is theoretically expected. This connection can be realized through nonperturbative Q^2 -dependence only. In my talk I discuss such nonperturbative Q^2 -dependence of the sum rules (see also [5]).

Below I will consider only the first moment of the structure function $g_1(x, Q^2)$

$$\Gamma_{p,n}(Q^2) = \int_0^1 dx \, g_{1;p,n}(x, Q^2) \quad (1)$$

The presentation of the material is divided into two parts. The first part deals with the case of high Q^2 . I discuss the determination of twist-4 contributions to $\Gamma_{p,n}$ by QCD sum rules and with the account of twist-4 corrections and of the uncertainties in their values compare the theory with the experiment. In the second part the case of low and intermediate $Q^2 \lesssim 1 \text{ GeV}^2$ is considered in the framework of the model, which realizes the smooth connection of GDH sum rule at $Q^2 = 0$ with the asymptotic form of $\Gamma_{p,n}(Q^2)$ at high Q^2 .

2. High Q^2

At high Q^2 with the account of twist-4 contributions $\Gamma_{p,n}(Q^2)$ have the form

$$\Gamma_{p,n}(Q^2) = \Gamma_{p,n}^{as}(Q^2) + \Gamma_{p,n}^{tw4}(Q^2) \quad (2)$$

$$\begin{aligned} \Gamma_{p,n}^{as}(Q^2) = \frac{1}{12} \left\{ [1 - a - 3.58a^2 - 20.2a^3 - ca^4] [\pm g_A + \frac{1}{3}a_8] \right. \\ \left. + \frac{4}{3} [1 - \frac{1}{3}a - 0.55a^2 - 4.45a^3] \Sigma \right\} - \frac{N_f}{18\pi} \alpha_s(Q^2) \Delta g(Q^2) \end{aligned} \quad (3)$$

$$\Gamma_{p,n}^{tw4}(Q^2) = \frac{b_{p,n}}{Q^2} \quad (4)$$

In eq.(3) $a = \alpha_s(Q^2)/\pi$, g_A is the β -decay axial coupling constant, $g_A = 1.260 \pm 0.002$ [6]

$$g_A = \Delta u - \Delta d \quad a_8 = \Delta u + \Delta d - 2\Delta s \quad \Sigma = \Delta u + \Delta d + \Delta s. \quad (5)$$

$\Delta u, \Delta d, \Delta s, \Delta g$ are parts of the nucleon spin projections carried by u, d, s quarks and gluons:

$$\Delta q = \int_0^1 \left[q_+(x) - q_-(x) \right] dx \quad (6)$$

where $q_+(x), q_-(x)$ are quark distributions with spin projection parallel (antiparallel) to nucleon spin and a similar definition takes place for Δg . The coefficients of perturbative series were calculated in [7-10], the numerical values in (3) correspond to the number of flavours $N_f = 3$, the coefficient c was estimated in [11], $c \approx 130$. In the $\overline{MS}$ renormalization scheme chosen in [7-10] a_8 and Σ are Q^2 -independent. In the assumption of the exact $SU(3)$ flavour symmetry of the octet axial current matrix elements over baryon octet states $a_8 = 3F - D = 0.59 \pm 0.02$ [12].

Strictly speaking, in (3) the separation of terms proportional to Σ and Δg is arbitrary, since the operator product expansion (OPE) has only one singlet in flavour twist-2 operator for the first moment of the polarized structure function – the operator of singlet axial current $j_{\mu 5}^{(0)}(x) = \sum_q \bar{q}_i(x) \gamma_\mu \gamma_5 q$, $q = u, d, s$. The separation of terms proportional to Σ and Δg is outside the framework of OPE and depends on the infrared cut-off. The expression used in (3) is based on the physical assumption that the virtualities p^2 of gluons in the nucleon are much larger than light quark mass squares, $|p^2| \gg m_q^2$ [13] and that the infrared cut-off is chosen in a way providing the standard form of axial anomaly [14].

Twist-4 corrections to $\Gamma_{p,n}$ were calculated by Balitsky, Braun and Koleshichenko (BBK) [15] using the QCD sum rule method.

BBK calculations were critically analyzed in [5], where it was shown that there are few possible uncertainties in these calculations: 1) the main contribution to QCD sum rules comes from the last accounted term in OPE – the operator of dimension 8; 2) there is a large background term and a much stronger influence of the continuum threshold comparing with usual QCD sum rules; 3) in the singlet case, when determining the induced by external field vacuum condensates, the corresponding sum rule was saturated by η -meson, what is wrong. The next order term – the contribution of the dimension 10 operator in the BBK sum rules was estimated by Oganessian [16]. The account of the dimension-10 contribution to the BBK sum rules and estimation of other uncertainties results in (see [5]):

$$b_{p-n} = -0.006 \pm 0.012 \text{ GeV}^2 \quad (7)$$

$$b_{p+n} = -0.035(\pm 100\%) \text{ GeV}^2 \quad (8)$$

As is seen from (7), in the nonsinglet case the twist-4 correction is small ($\lesssim 2\%$ at $Q^2 \gtrsim 5 \text{ GeV}^2$) even with the account of the error. In the singlet case the situation is much worse: the estimate (8) may be considered only as correct by the order of magnitude.

I turn now to comparison of the theory with the recent experimental data. In Table 1 the recent data obtained by SMC [1] and E 154(SLAC) [2] groups are presented.

Table 1

	Γ_p	Γ_n	$\Gamma_p - \Gamma_n$	$\alpha_s(5GeV^2)$
SMC	0.132 ± 0.017	-0.048 ± 0.022	0.181 ± 0.035	$0.270^{+0.16}_{-0.40}$
combined	0.142 ± 0.011	-0.061 ± 0.016	0.202 ± 0.022	$0.116^{+0.16}_{-0.44}$
E 154(SLAC)	0.112 ± 0.014	-0.056 ± 0.008	0.168 ± 0.012	$0.339^{+0.052}_{-0.063}$
EJ/Bj sum rules	0.168 ± 0.005	-0.013 ± 0.005	0.181 ± 0.002	0.276

In the second line of Table 1 the results of the performed by SMC [1] combined analysis of SMC [1], SLAC-E80/130 [17], EMC [18] and SLAC-E143 [19] data are given. The data presented in Table 1 refer to $Q^2 = 5GeV^2$. In all measurements each range of x corresponds to each own mean Q^2 . Therefore, in order to obtain $g_1(x, Q^2)$ at fixed Q^2 refs. [1,2] use the following procedure. At some reference scale Q_0^2 ($Q_0^2 = 1GeV^2$ in [1] and $Q_0^2 = 0.34GeV^2$ in [2]) quark and gluon distribution were parametrized as functions of x . (The number of the parameters was 12 in [1] and 8 in [2]). Then NLO evolution equations were solved and the values of the parameters were determined from the best fit at all data points. The numerical values presented in Table 1 correspond to $\overline{MS}$ regularization scheme, statistical, systematical, as well as theoretical errors arising from uncertainty of α_s in the evolution equations, are added in quadratures. In the last line of Table 1 the Ellis-Jaffe (EJ) and Bjorken (Bj) sum rules prediction for Γ_p, Γ_n and $\Gamma_p - \Gamma_n$, correspondingly are given. The EJ sum rule prediction was calculated according to (3), where $\Delta s = 0$, i.e., $\Sigma = a_8 = 0.59$ was put and the last-gluonic term in (3) was omitted. The twist-4 contribution was accounted in the Bj sum rule and included into the error in the EJ sum rule. The α_s value in the EJ and Bj sum rules calculation was chosen as $\alpha_s(5GeV^2) = 0.276$, corresponding to $\alpha_s(M_z) = 0.117$ and $\Lambda_{\overline{MS}}^{(3)} = 360MeV$ (in two loops). As is clear from Table 1, the data, especially for Γ_n , contradict the EJ sum rule. In the last column, the values of α_s determined from the Bj sum rule are given with the account of twist-4 corrections.

The experimental data on Γ_p presented in Table 1 are not in a good agreement. Particularly, the value of Γ_p given by E154 Collaboration seems to be low: it does not agree with the old data presented by SMC [20] ($\Gamma_p = 0.136 \pm 0.015$) and E143 [19] ($\Gamma_p = 0.127 \pm 0.011$). Even more strong discrepancy is seen in the values of α_s , determined from the Bj sum rules. The value which follows from the combined analysis is unacceptably low: the central point corresponds to $\Lambda_{\overline{MS}}^{(3)} = 15MeV$! On the other side, the value, determined from the E154 data seems to be high, the corresponding $\alpha_s(M_z) = 0.126 \pm 0.009$. Therefore, I come to a conclusion that at the present level of experimental accuracy α_s cannot be reliably determined from the Bj sum rule in polarized scattering.

Table 2 shows the values of Σ – the total nucleon spin projection carried by u, d and s -quarks found from Γ_p and Γ_n presented in Table 1 using eq.(3). (It was put $g_A = 1.260, a_8 = 0.59$, the term, proportional to Δg is included into Σ).

Table 2: The values of Σ

	From Γ_p		From Γ_n	
	At $\alpha_s(5GeV^2) = 0.276$	At $\alpha_s(GeV^2)$ given in Table 1	At $\alpha_s(5GeV^2) = 0.276$	At $\alpha_s(5GeV^2)$ given in Table 1
SMC	0.256	0.254	0.264	0.266
Comb.	0.350	0.250	0.145	0.225
E154	0.070(0.14; 0.25)	0.133(0.20; 0.30)	0.19(0.25; 0.14)	0.144 (0.205; 0.10)

In their fitting procedure [2] E154 Collaboration used the values $a_8 = 0.30$ and $g_A = 1.09$. The values of Σ obtained from Γ_p and Γ_n given by E154 at $a_8 = 0.30, g_A = 1.26$ and $a_8 = 0.30, g_A = 1.09$ are presented in parenthesis. The value $a_8 = 0.30$ corresponds to a strong violation of SU(3) flavour symmetry and is unpalusible; $g_A = 1.09$ means a bad violation of isospin and is unacceptable. As seen from Table 1, Σ is seriously affected by these assumptions. The values of Σ found from Γ_p and Γ_n using SMC and combined analysis data agree with each other only, if one takes for $\alpha_s(5GeV^2)$ the values given in Table 1 ($\alpha_s = 0.116$ for combined data), what is unacceptable. The twist-4 corrections were not accounted in Σ in Table 2: their account, using eqs.(7), (8), results in increasing of Σ by 0.04 if determined from Γ_p and by 0.03 if determined from Γ_n .

To conclude, one may say, that the most probable value of Σ is $\Sigma \approx 0.3$ with an uncertain error. The contribution of gluons may be estimated as $\Delta g(1GeV^2) \approx 0.3$ (see [5], [21], [22]). Then $\Delta g(5GeV^2) \approx 0.6$ and the account of gluonic term in eq.(3) results in increasing of Σ by 0.06. At $\Sigma = 0.3$ we have $\Delta u = 0.83, \Delta d = -0.43, \Delta s = -0.1$.

3. Low and Intermediate Q^2

The problem of a smooth connection of the Gerasimov-Drell-Hearn (GDH) sum rules [3,4] which holds at $Q^2 = 0$, and the sum rules at high Q^2 attracts attention in the last years [5, 23-25].

In order to connect the GDH sum rule with $\Gamma_{p,n}(Q^2)$ consider the integrals [26]

$$I_{p,n}(Q^2) = \int_{Q^2/2}^{\infty} \frac{d\nu}{\nu} G_{1;p,n}(\nu, Q^2) \quad (9)$$

Changing the integration variable ν to x , (9) can be also identically written as

$$I_{p,n}(Q^2) = \frac{2m^2}{Q^2} \int_0^1 dx g_{1;p,n}(x, Q^2) = \frac{2m^2}{Q^2} \Gamma_{p,n}(Q^2) \quad (10)$$

At $Q^2 = 0$ the GDH sum rule takes place

$$I_p(0) = -\frac{1}{4}\kappa_p^2 = -0.8035; \quad I_n(0) = -\frac{1}{4}\kappa_n^2 = 0.9149; \quad I_p(0) - I_n(0) = 0.1114 \quad (11)$$

where κ_p and κ_n are proton and neutron anomalous magnetic moments.

The schematic Q^2 dependence of $I_p(Q^2)$, $I_n(Q^2)$ and $I_p(Q^2) - I_n(Q^2)$ is plotted in Fig.1. The case of $I_p(Q^2)$ is especially interesting: $I_p(Q^2)$ is positive, small and decreasing at $Q^2 \gtrsim 3GeV^2$ and negative and relatively large in absolute value at $Q^2 = 0$. With

$I_n(Q^2)$ the situation is similar. All this indicates large nonperturbative effects in $I(Q^2)$ at $Q^2 \lesssim 1\text{GeV}^2$.

In [23] the model was suggested, which describes $I(Q^2)$ (and $\Gamma(Q^2)$) at low and intermediate Q^2 , where GDH sum rules and the behaviour of $I(Q^2)$ at large Q^2 were fulfilled. The model had been improved in [24]. (Another model with the same goal was suggested by Soffer and Teryaev [25]).

Since it is known, that at small Q^2 the contribution of resonances to $I(Q^2)$ is of importance, it is convenient to represent $I(Q^2)$ as a sum of two terms

$$I(Q^2) = I^{res}(Q^2) + I'(Q^2), \quad (12)$$

where $I^{res}(Q^2)$ is the contribution of baryonic resonances. $I^{res}(Q^2)$ can be calculated from the data on electroproduction of resonances. Such calculation was done with the account of resonances up to the mass $W = 1.8\text{GeV}$ [27].

In order to construct the model for nonresonant part $I'(Q^2)$ consider the analytical properties of $I(Q^2)$ in q^2 . As is clear from (9),(10), $I(Q^2)$ is the moment of the structure function, i.e. it is a vertex function with two legs, corresponding to ingoing and outgoing photons and one leg with zero momentum. The most convenient way to study of analytical properties of $I(q^2)$ is to consider a more general vertex function $I(q_1^2, q_2^2; p^2)$, where the momenta of the photons are different, and go to the limit $p \rightarrow 0$, $q_1^2 \rightarrow q_2^2 = q^2$. $I(q^2, q_2^2; p^2)$ can be represented by the double dispersion relation:

$$I(q^2) = \lim_{q_1^2 \rightarrow q_2^2 = q^2, p^2 \rightarrow 0} I(q_1^2, q_2^2; p^2) = \left\{ \int ds_2 \int ds_1 \frac{\rho(s_1, s_2; p^2)}{(s_1 - q_1^2)(s_2 - q_2^2)} + \right. \\ \left. + P(q_1^2) \int \frac{\varphi(s, p^2)}{s - q_2^2} ds + P(q_2^2) \int \frac{\varphi(s, p^2)}{s - q_1^2} ds \right\}_{q_1^2 = q_2^2 = q^2, p \rightarrow 0} \quad (13)$$

The last two terms in (13) are the subtraction terms in the double dispersion relation, $P(q^2)$ is the polynomial. According to (10), $I(q^2)$ decreases at $|q^2| \rightarrow \infty$, $P(q^2) = \text{Const}$ and the constant subtraction term in (13) is absent. We are interesting in $I(Q^2)$ dependence in the domain $Q^2 \lesssim 1\text{GeV}^2$. Since after performed subtraction, the integrals in (13) are well converging, one may assume, that at $Q^2 \lesssim 2-3\text{GeV}^2$ the main contribution comes from vector meson intermediate states, so the general form of $I'(Q^2)$ is

$$I'(Q^2) = \frac{A}{(Q^2 + \mu^2)^2} + \frac{B}{Q^2 + \mu^2}, \quad (14)$$

where A and B are constants, μ is ρ (or ω) mass. The constants A and B are determined from GDH sum rules at $Q^2 = 0$ and from the requirement that at high $Q^2 \gg \mu^2$ takes place the relation

$$I(Q^2) \approx I'(Q^2) \approx \frac{2m^2}{Q^2} \Gamma^{as}(Q^2), \quad (15)$$

where $\Gamma^{as}(Q^2)$ is given by (3). ($I^{res}(Q^2)$ fastly decreases with Q^2 and is very small above $Q^2 = 3\text{GeV}^2$). These conditions are sufficient to determine in unique way the constant A and B in (14). For $I'(Q^2)$ it follows:

$$I'(Q^2) = 2m^2\Gamma^{as}(Q_m^2)\left[\frac{1}{Q^2 + \mu^2} - \frac{c\mu^2}{(Q^2 + \mu^2)^2}\right], \quad (16)$$

$$c = 1 + \frac{\mu^2}{2m^2} \frac{1}{\Gamma^{as}(Q_m^2)} \left[\frac{1}{4}\kappa^2 + I^{res}(0)\right], \quad (17)$$

where $I_p^{res}(0) = -1.03$, $I_n^{res}(0) = -0.83$ [24].

The model and eq.16 cannot be used at high $Q^2 \gtrsim 5GeV^2$: one cannot believe, that at such Q^2 the saturation of the dispersion relation (13) by the lowest vector meson is a good approximation. For this reason there is no matching of (16) with QCD sum rule calculations of twist-4 terms. (Formally, from (16) it would follow $b_{p-n} \approx -0.15$, $b_{p+n} \approx -0.07$). It is not certain, what value of the matching point Q_m^2 should be chosen in (16). This results in 10% uncertainty in the theoretical predictions. Fig.'s 2,3 shows the predictions of the model in comparison with recent SLAC data [28], obtained at low $Q^2 = 0.5$ and $1.2GeV^2$ as well as SMC and SLAC data at higher Q^2 . The chosen parameters are $\Gamma_p(Q_m^2) = 0.142$, $\Gamma_n(Q_m^2) = -0.061$, corresponding to $c_p = 0.458$, $c_n = 0.527$ in (16),(17). The agreement with the data, particularly at low Q^2 , is very good. The change of the parameters only weakly influences $\Gamma_{p,n}(Q^2)$ at low Q^2 .

This work was supported in part by the Russian Foundation of Fundamental Research Grant 97-02-16131, CRDF Grant RP2-132 and Schweizerische Nationalfond Grant 7SUPJ048716.

References

- [1] SMC-Collaboration, hep-ph/9702005, submitted to Physical Review D.
- [2] K.Abe et al (E154 Collab.) Phys.Lett. **B405** (1997) 180.
- [3] S.Gerasimov, Yad.Fiz. **2** (1996) 930.
- [4] S.D.Drell and A.C.Hearn, Phys.Rev.Lett. **16** (1996) 908.
- [5] B.L.Ioffe, hep-ph/9704295, Phys.At.Nucl. in press.
- [6] R.M.Barnett et al., Particle Data Group, Phys.Rev. **D54** (1996) 1.
- [7] J.Kadaira et al., Phys.Rev. **D20** (1979) 627; Nucl.Phys. **B159** (1979) 99, **165** (1980) 129.
- [8] S.A.Larin and J.A.M.Vermaseren, Phys.Lett. **259** (1991) 345.
- [9] S.A.Larin, Phys.Lett. **334** (1994) 192.
- [10] S.A.Larin, T.van Ritbergen and J.A.M.Vermaseren, preprint NIKHEF-97-011 (1997).
- [11] A.L.Kataev, Phys.Rev. **D50** (1994) 5469.
- [12] S.Y.Hsueh et al., Phys.Rev. **D38** (1988) 2056.
- [13] R.D.Carlitz, J.C.Collins and A.H.Mueller, Phys.Lett. **B 214** (1988) 229.
- [14] S.D.Bass, B.L.Ioffe, N.N.Nikolaev and A.W.Thomas, J.Moscow Phys.Soc. **1** (1991) 317.
- [15] I.I.Balitsky, V.M.Braun and A.V.Kolesnichenko, Phys.Lett. **242** (1990) 245, Errata **B318** (1993) 648.
- [16] A.Oganesian, hep-ph/9704435, Phys.At.Nucl., in press.
- [17] M.J.Alguard et al, Phys.Rev.Lett. **37** (1976) 1261; **41** (1978) 70. G.Baum, ibid **51** (1983) 1135.
- [18] J.Ashman et al, Phys.Lett **206** (1988) 364; Nucl.Phys. **B328** (1989) 1.
- [19] K.Abe et al. (SLAC E143 Collaboration), Phys.Rev.Lett. **74** (1995) 346.
- [20] D.Adams et al, Phys.Lett. **B329** (1994) 399.
- [21] I.Balitsky, X.Ji, Phys.Rev.Lett. **79** (1997) 1225.
- [22] A.Saalfeld, G.Piller, L.Mankiewicz, Preprint TUM/T39-97-19, hep-ph/9708378.
- [23] M.Anselmino, B.L.Ioffe and E.Leader, Sov.J.Nucl.Phys. **49** (1989) 136.
- [24] V.Burkert and B.L.Ioffe, JETP, **105** (1994) 1153.

- [25] J.Soffer and O.Teryaev, Phys.Rev. **D51** (1995) 25.
- [26] B.L.Ioffe, V.A.Khoze, L.N.Lipatov, Hard Processes, v.1, North Holland, Amsterdam, 1984.
- [27] V.Burkert and Z.Li, Phys.Rev. **47** (1993) 46.
- [28] K.Abe et al., (SLAC E143 Collaboration) Phys.Rev.Lett. **78** (1997) 815.
- [29] P.L.Anthony et al. (SLAC E142 Collaboration),
Phys.Rev. **D54** (1996) 6620.
- [30] K.Abe et al. (SLAC E143 Collaboration), Phys.Rev.Lett. **75** (1995) 25.

Figure Captions

- Fig. 1** The Q^2 -dependence of integrals $I_p(Q^2), I_n(Q^2), I_p(Q^2) - I_n(Q^2)$. The vertical axis is broken at negative values.
- Fig. 2** The Q^2 -dependence of $\Gamma_p = \Gamma'_p + \Gamma_p^{res}$ (solid line), described by eqs.(12,16,17). Γ_p^{res} (dotted) and Γ'_p (dashed) are the resonance and nonresonance parts. The experimental points are: the dots from E143 [28], the square - from E143 (SLAC) [19], the cross - SMC-SLAC combined data [1], the triangle from SMC [1].
- Fig. 3** The same as in Fig.2, but for neutron. The experimental points are: the dots from E143 (SLAC) measurements on deuteron [28], the square at $Q^2 = 2GeV^2$ is the E142(SLAC) [29] data from measurements on polarized 3He , the square at $Q^2 = 3GeV^2$ is E143(SLAC) [30] deuteron data, the cross is SMC-SLAC combined data [1], the triangle is SMC deuteron data [1].




Figure 1:




Figure 2:




Figure 3: